

RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

FIFRA Section 24(c) SLN LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

rozol[®] VOLE BAIT

FOR CONTROL OF MEADOW OR MOUNTAIN VOLES IN POME FRUIT (APPLE, PEAR) AND STONE FRUIT (APRICOT, CHERRY, NECTARINE, PEACH, PLUM, PRUNE) TREE ORCHARDS

EPA SLN No. WA-070019

EPA Reg. No. 7173-242

This label for Rozol Vole Bait expires and must not be distributed or used in accordance with this SLN registration after December 31, 2017.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions and precautions on the EPA registered label.

DIRECTIONS FOR USE:

POME AND STONE FRUIT TREE ORCHARDS FOR MEADOW OR MOUNTAIN VOLE CONTROL:

APPLICATION DIRECTIONS: Before application, examine orchard floor to locate trails and runway systems to be treated. Apply bait when three days of rain-free and snow-free weather are expected.

Hand spot baiting: Place about 1-1/2 ounces of bait in each active hole, trail or runway at each tree site. Do not exceed 10 pounds of bait per acre. Cover each placement of bait to avoid exposing nontarget organisms.

Ground or aerial broadcast baiting: Using a commercial spreader, uniformly apply 10 pounds of bait per acre. If populations are high, make a second application 1 to 2 months after the first, not to exceed 10 pounds of bait per acre. Do not apply more than 20 pounds of bait per acre per season. To avoid exposing nontarget organisms, clean up any bait in piles for reuse or disposal according to the federal label.

RESTRICTIONS / PRECAUTIONS:

- Do not apply more than 20 pounds of bait per acre per season.
- Apply bait after harvest (including all harvestable fruit drops) and before new growth in the spring. Do not apply bait during the growing season.
- Do not apply bait over bare ground or to crops not specified above.
- Do not expose humans (including children), pets, livestock or other nontarget animals to bait during application, handling, storage or disposal. Do not allow livestock to graze or clean up fruit drops in treated areas. Do not use hay cut after application for livestock feed or bedding.
- This pesticide is toxic to fish and wildlife. Rozol Vole Bait should not be used under this SLN label where impact on listed threatened or endangered species is likely. Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply by air within 50 feet of surface water (i.e. streams, rivers, lakes). You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

WSDA Container Disposal Guidance:

Completely empty the contents of the container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington. (020613)

24(c) registrant:
Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209

155-0009-0213
Rev. 02/06/13